



L.A. Care
HEALTH PLAN®

For All of L.A.

2018 HEDIS Cal MediConnect				
Acronyms	Measures	Rate 2016 HEDIS (CY 2015)	Rate 2017 HEDIS (CY 2016)	Rate 2018 HEDIS (CY 2017)
AAP	Adults Access to Preventive/Ambulatory Health Services	75.4%	77.1%	79.5%
ABA	Adult BMI Assessment	87.1%	93.9%	95.8%
AMM	Antidepressant Medication Management – Acute Phase	48.3%	64.2%	65.7%
AMM	Antidepressant Medication Management – Continuous Phase	34.6%	46.3%	53.9%
ART	Rheumatoid Arthritis Management	71.0%	73.9%	72.0%
BCS	Breast Cancer Screening	61.2%	62.6%	60.1%
COL	Colorectal Cancer Screening	45.3%	48.4%	57.7%
CBP	Controlling High Blood Pressure	56.2%	66.9%	69.5%
COA	Care for Older Adults – Medication Review	58.4%	64.2%	NQ
COA	Care for Older Adults – Functional Status Assessment	38.4%	41.1%	NQ
COA	Care for Older Adults – Pain Screening	57.9%	62.0%	NQ
CDC	Diabetes Care - Eye Exams	65.0%	64.2%	70.4%
CDC	Diabetes Care – Kidney Disease	95.0%	95.9%	96.8%
CDC	Diabetes Care – Blood Sugar Controlled	42.3%	56.5%	62.5%
DAE	Use of High Risk Medications in the Elderly – at least one drug	19.6%	13.5%	25.0%

NQ-Not required to report



L.A. Care
HEALTH PLAN®

For All of L.A.

2018 HEDIS Cal MediConnect				
Acronyms	Measures	Rate 2016 HEDIS (CY 2015)	Rate 2017 HEDIS (CY 2016)	Rate 2018 HEDIS (CY 2017)
DAE	Use of High Risk Medications in the Elderly – at least two drugs	3.7%	7.8%	12.4%
DDE	Potentially Harmful Drug-Disease Interactions in the Elderly – Total	47.3%	39.8%	45.1%
FUH	Follow-Up after Hospitalization for Mental Illness –within 30-days	11.9%	41.9%	46.9%
FUH	Follow-Up after Hospitalization for Mental Illness -within 7-days	7.7%	25.9%	28.1%
IET	Initiation and Engagement of Alcohol and Other Drug Dependence Treatment - Engagement	2.5%	NA	3.3%
IET	Initiation and Engagement of Alcohol and Other Drug Dependence Treatment – Initiation	34.0%	32.5%	38.9%
PBH	Persistence of Beta Blocker Treatment After a Heart Attack	86.1%	85.4%	88.2%
PCE	Pharmacotherapy Management of COPD Exacerbation- Bronchodilators	89.6%	82.75%	85.1%
PCE	Pharmacotherapy Management of COPD Exacerbation -Systemic Corticosteroid	63.2%	60.9%	61.2%
OMW	Osteoporosis Management in Women Who Had a Fracture	29.0%	31.7%	27.3%
SPR	Use of Spirometry Testing in the Assessment and Diagnosis of COPD	16.6%	16.3%	17.6%

NQ-Not required to report