



2019 HEDIS Cal MediConnect				
Acronyms	Measures	Rate 2017 HEDIS (CY 2016)	Rate 2018 HEDIS (CY 2017)	Rate 2019 HEDIS (CY 2018)
AAP	Adults Access to Preventive/Ambulatory Health Services	77.1%	79.5%	82.4%
ABA	Adult BMI Assessment	93.9%	95.8%	97.1%
AMM	Antidepressant Medication Management – Acute Phase	64.2%	65.7%	64.8%
AMM	Antidepressant Medication Management – Continuous Phase	46.3%	53.9%	57.2%
ART	Rheumatoid Arthritis Management	73.9%	72.0%	75.7%
BCS	Breast Cancer Screening	62.6%	60.1%	63.7%
COL	Colorectal Cancer Screening	48.4%	57.7%	61.0%
CBP	Controlling High Blood Pressure	66.9%	69.5%	73.2%
COA	Care for Older Adults – Medication Review	64.2%	NQ	71.8%
COA	Care for Older Adults – Functional Status Assessment	41.1%	NQ	52.8%
COA	Care for Older Adults – Pain Screening	62.0%	NQ	74.7%
CDC	Diabetes Care - Eye Exams	64.2%	70.4%	75.6%
CDC	Diabetes Care – Kidney Disease	95.9%	96.8%	97.0%
CDC	Diabetes Care – Blood Sugar Controlled	56.5%	62.5%	62.2%
DAE	Use of High Risk Medications in the Elderly – at least one drug	13.5%	25.0%	25.8%

NQ-Not required to report



L.A. Care
HEALTH PLANSM

For All of L.A.

2019 HEDIS Cal MediConnect				
Acronyms	Measures	Rate 2017 HEDIS (CY 2016)	Rate 2018 HEDIS (CY 2017)	Rate 2019 HEDIS (CY 2018)
DAE	Use of High Risk Medications in the Elderly – at least two drugs	7.8%	12.4%	13.0%
DDE	Potentially Harmful Drug-Disease Interactions in the Elderly – Total	39.8%	45.1%	40.5%
FUH	Follow-Up after Hospitalization for Mental Illness –within 30-days	41.9%	46.9%	49.0%
FUH	Follow-Up after Hospitalization for Mental Illness -within 7-days	25.9%	28.1%	29.5%
IET	Initiation and Engagement of Alcohol and Other Drug Dependence Treatment - Engagement	NA	3.3%	4.6%
IET	Initiation and Engagement of Alcohol and Other Drug Dependence Treatment – Initiation	32.5%	38.9%	42.2%
PBH	Persistence of Beta Blocker Treatment After a Heart Attack	85.4%	88.2%	*
PCE	Pharmacotherapy Management of COPD Exacerbation- Bronchodilators	82.75%	85.1%	83.3%
PCE	Pharmacotherapy Management of COPD Exacerbation -Systemic Corticosteroid	60.9%	61.2%	76.0%
OMW	Osteoporosis Management in Women Who Had a Fracture	31.7%	27.3%	29.4%
SPR	Use of Spirometry Testing in the Assessment and Diagnosis of COPD	16.3%	17.6%	20.3%

*Denominator less than 30

NQ-Not required to report