WELCOME!

Before we get started:

- Please scan the QR code or visit the following link to register
 - https://forms.gle/AWiMBWmMWajdEkwJA
 - The QR code/link should bring you to the following page (PLEASE FILL OUT ALL MANDATORY QUESTIONS):







Early detection. Better care.



Learning Objectives

- 1. UNDERSTAND THE WHY'S AND WHAT'S OF DEMENTIA SCREENING AND CARE
- 2. LEARN HOW TO CONDUCT A SCREENING CALLED THE COGNITIVE HEALTH ASSESSMENT: TOOLS, SCORING, AND INTERPRETING

Step 1: Take a Brief Patient History

Step 2: Use Screening Tools

Step 3: Document Care Partner Information

- 1. THERE'S A POSITIVE SCREEN, WHAT ARE THE NEXT STEPS?
- 2. BILLING REQUIREMENTS



The Impact of Dementia is Increasing in CA

BY THE NUMBERS = Prevalence, Lifetime Risks, Cause of Death

- Between 2019 and 2040, the **population of California will expand by 16%**, whereas the population of people living with Alzheimer's disease (AD) **will expand by 127%**.
- Among California's residents who live to be 65, one in five people will develop dementia.
- Between 2014 and 2017, AD accounted for 28% as the cause of death (the greatest increase in causes of death in

(Ross, L, Beld, M, and Yeh, J. (2021). Alzheimer's Disease and Related Dementias Facts and Figures in California: Current Status and Future Projections. Report prepared for the California Department of Public Health, Sacramento, CA at the Institute for Health and Aging, University of California, San Francisco, CA.)



The Types of Cognitive Decline: More Than Just Memory



*Cognitive decline **is not** just memory decline.*



The Types of Cognitive Decline: Six Domains of Cognition

- 1. The Learning and Memory Domain
- 1. Executive Function
- 2. Complex Attention
- 3. Visuospatial Function
- **4. Social Cognition**
- 5. Language





Types of Cognitive Decline

Type of Cognitive Decline	Magnitude of decline	Affects Daily Function?
Age-related decline	Normal decline in cognitive functions for age	No
Mild Cognitive Impairment (MCI)	Abnormal decline in cognitive functions for age	No. May be using compensatory strategies to accomplish activities of daily living.
Dementia	Abnormal decline in cognitive functions for age	Yes. Unable to use compensatory strategies to accomplish activities of daily living.



Suggests Age-Related Cognitive Decline

Suggests Abnormal Cognitive Decline

The patient cannot remember details of a conversation or event that took place a year ago.

The patient cannot recall details of recent events or conversations.

The patient cannot remember the name of an acquaintance they don't see frequently.

The patient cannot remember the names of close family members.

The patient forgets things and events occasionally.

The patient forgets things or events nearly on a daily basis.

The patient occasionally has difficulty finding words.

The patient frequently pauses and substitutes when finding words.

The patient is worried about their memory, but their relatives are not.

The patient's relatives are worried about their memory, but the patient is not aware of any problems.

REMEMBER:

Age-Related Cognitive Decline Is NOT a Disease

What Cognitive Functions Decline in Normal Aging?

COGNITIVE FUNCTIONS THAT ARE VULNERABLE TO THE DECLINE IN AGING ARE:

- Short-term memory
- More time and effort to recall new information
- Decreased efficiency (e.g., divided attention and multitasking)
- Slower learning speed



Did You Know?

It's often someone who knows the patient well who first notices signs or

symptoms of cognitive decline and brings them to the attention of a medical

provider. However, many also attribute these signs to "old age" and may not

think of bringing them up to a provider.



What Is Mild Cognitive Impairment (MCI) or Mild Neurocognitive Disorder (MINCD)?

A.When a person shows subjective signs or symptoms of cognitive decline in one or more cognitive domains

A.These changes are substantiated by cognitive testing.

B.The person has an intact ability to live independently and shows no impairment in social and/or occupational functioning.



What Is Dementia?

Dementia is also known as Major Neurocognitive Disorder (MaNCD).

Persons must exhibit the following three criteria:

- 1. Acquired Cognitive Decline
- 2. Acquired Functional Decline
- 3. No Other Causes



Acquired Cognitive Decline

1. The person must have an acquired cognitive decline from their prior level of ability in at least one cognitive domain.

Remember the SIX Domains?

learning and memory, language, executive function, complex attention, visuospatial skills, and social cognition.




Acquired Decline in Function

- 2. The person must have an acquired decline in function from their prior level of ability:
 - One or more Instrumental Activities of Daily Living (IADLs) or Activities of Daily Living (ADLs).



No Other Causes

The patient must not have any other medical or psychiatric disorders that may explain their cognitive decline.






DEMENTIA: AN UMBRELLA TERM

Dementia is an "umbrella" term that describes the syndrome of cognitive and functional decline.

Alzheimer's disease, the most common, makes up 50%-75% of all cases.

Vascular dementia is thought to cause 20%-30% of cases and often coexists with AD, also called "mixed dementia."







t's REVIEW and try some case stud

ies.



MEET MR. AHOKA: Please read case

Mr. Ahoka is a **71-year-old man** who has poorly controlled hypertension, had a stroke 20 years ago, and uses tobacco.

Several family members have dementia.

He reports he has trouble "recalling conversations" and misplaces things. He lost his phone twice. He gets very angry when he can't find things, and this was not his personality before.





Let's Review Mr. Ahoka's Screening Results

- Mr. Ahoka's score on a validated brief cognitive screen suggested cognitive impairment.
- His PCP then reviewed his **functional abilities** by asking about basic and instrumental activities of daily living. Mr. Ahoka reported that he cannot independently do instrumental activities of daily living. For example, his sister now pays his bills, does his shopping, and prepares his meals.



At a follow-up after this positive cognitive health assessment screening result:

- His PCP ordered lab work searching for reversible causes of cognitive impairment and reviewed his medications for possible cognitive side effects, and found **no clear causes** for his symptoms.
- Mr. Ahoka also had a negative screen for depression and substance use.



Which type of cognitive impairment do you think Mr. Ahoka is showing?

1.Age-Related Decline

2. Mild Cognitive Impairment (MCI)

3. Dementia



MEET MS. HADBURY

- Ms. Hadbury is a 73-year-old woman whose mother had late-life dementia.
- She reports a concern for cognitive decline to her PCP. The patient has insight into her cognitive symptoms and provides several examples to illustrate what she experiences.
- She cannot pick up new skills or information as well, such as a recent class she took in which she could not remember the information from one class to the next.
- She also noticed that drawing, a hobby she loves, has become more difficult.





Let's Review Ms. Hadbury's Screening Results

- Ms. Hadbury's score on a validated brief cognitive screen suggests cognitive impairment.
- Her PCP then reviewed her functional abilities by asking about basic and instrumental activities of daily living. Ms. Hadbury says in the last two years she has had to take a list with her to the grocery store or she'll forget what she came for, but otherwise **she is doing all activities independently.**



After this positive cognitive health assessment screening result:

- Her PCP ordered lab work searching for reversible causes of cognitive impairment, and all results are normal. She's not on any medications that have cognitive side effects.
- Ms. Hadbury also had a negative screen for depression and substance use



Which type of cognitive impairment do you think Ms. Hadbury is showing?

1.Age-Related Decline

2. Mild Cognitive Impairment (MCI)

3. Dementia



CONGRATULATIONS! Now you know the three types of cognitive decline. Let's discuss the benefits of early dementia detection.

Benefits of Early Cognitive Assessments

- Improves quality of life for affected persons and caregivers.
- Reduces unnecessary costs of care.
- Increases the likelihood of benefiting from evidence-based dementia prevention strategies.
- Can address racial disparities in cognitive impairment among Black, Indigenous, Latino and other communities which are currently underserved.

Introducing the "Cognitive Health Assessment"



The CHA is Designed Specifically for Primary Care Providers and Busy Clinics

It includes assessments that are

- Free to use
- Quick to administer
- Easy to score
- Validated in primary care
- Available in multiple languages





A Starting Place

Learning the CHA will enable you to start the process of

detecting cognitive decline in your patient.





HOW TO CONDUCT THE CHA "Cognitive Health Assessment"



Three Key Steps

1 Take a brief patient history.

2 Use screening tools.

3 Document care partner information.








Start by Taking a Brief Patient History

History or signs of decline can come from many sources—from the patient, an informant, or you or your team members.

Remember to document the source of the information.





Obtaining and documenting a brief patient history is the foundation to start a brain health plan.

Tip: Establish a Partnership

If possible, involve a patient's support person as soon as possible, starting with history-taking. It sets the stage for an ongoing partnership in care—between you, the patient, and their support person. This also helps in the next step of the assessment.



Step 2: Use Screening Tools

The next CHA step is to assess

- the patient's cognition
- the patient's function and
- obtain collateral information from an informant.





The table below provides examples of validated tools for assessing a patient's cognition and function with the patient and an informant.

	Cognitive Screen Tools	Functional Screen Tools	
Patient	GP-COG OR Mini-Cog	ADL/IADL	
Informant	AD-8 OR Short IQ-CODE	GP-COG Informant Interview OR FAQ	_

GP-COG: General Practitioner assessment of Cognition

Mini-Cog: This is a short cognitive assessment; Mini-Cog is not a shortened name.

ADL: Activities of Daily Living

IADL: Instrumental Activities of Daily

Living

AD-8: Eight-item Informant Interview to Differentiate Aging and Dementia **Short IQ-CODE**: Short Informant Questionnaire

on Cognitive Decline in the Elderly

FAQ: Functional Activities

Questionnaire

GP-COG: Overview and Scoring

Quick facts:

- •Time to administer: 4-5 min
- Domains tested: visuospatial, executive, orientation, memory
- Accessibility: Available in multiple languages
- Web-based with automatic scoring
- Noteworthy: The address used in the test could be strange to people
- Not adjusted for a patient's education level

Scoring: There are a total of 9 points. The first question regarding name and address is worth up to 5 points. The other items are worth one point each. 0-4 indicates cognitive impairment. 5-8 indicates more information is

needed.

9 (out of 9) indicates no significant cognitive impairment.



GP-COG Informant Interview

Quick facts:

- •Time to administer: 2 min
- Six questions
- Web-based with automatic scoring
- Accessibility: Available in multiple languages

Scoring: "Yes" responses indicate impairment. Any question not answered with a "yes" counts as 1 point. There are a total of 6 points.

- 0-3 indicates cognitive impairment.
- 4-5 indicates less impairment.



Patient name:	Date:
Формунаровический повы поситованевы постаневы с	

GPCOG Screening Test

Step 1: Patient Examinatio Unless specified, each question should only be a		
Name and Address for subsequent recall test		
1. "1 am going to give you a name and address. After I have said it, it. Remember this name and address because I am going again in a few minutes: John Brown, 42 West Street, Ken of 4 attempts).	g to ask you to	tell it to me
Time Orientation	Correct I	ncorrect
2. What is the date? (exact only)		
Clock Drawing - use blank page		
3. Please mark in all the numbers to indicate the hours of a clock (correct spacing required)		
1. Please mark in hands to show 10 minutes past eleven o'clock (11.10)		
Information		
 Can you tell me something that happened in the news red (Recently = in the last week. If a general answer is given eg "war", "lot of rain", ask for details. Only specific answer so 	n, ,	
Recall		
1. What was the name and address I asked you to		
remember		
John -		
Brown	17	
42		
West (St)		
Kensington	44.	\neg
(To get a total score, add the number of items answered correct Total correct (score out of 9)	tly 19	9

<u>If patient scores 9, no significant cognitive impairment and further testing not necessary.</u>
<u>If patient scores 5-8, more information required. Proceed with Step 2, informant section.</u>
If patient scores 0-4, cognitive impairment is indicated. Conduct standard investigations.

University of New South Wales as represented by the Dementia Collaborative Research Centre — Assessment and Better Care; Brodaty et al, *JAGS* 2002; 50:530-534

Informant interview

		Date:		
Informant's name:				
In	formant's relationship to patient, i.e. informant is the patient	's:		
	These six questions ask how the patient is compared was well, say 5 — 10 years ago	d to when s/he		
	Compared to a few years ago:			
		Don't Yes No Know NIA		
	Does the patient have more trouble remembering things that have happened recently than s/he used to?			
	Does he or she have more trouble recalling conversations a few days later?			
	When speaking, does the patient have more difficulty in finding the right word or tend to use the wrong words more often?			
	 Is the patient less able to manage money and financial affairs (e.g. paying bills, budgeting)? 			
	• Is the patient less able to manage his or her medication independently?			
•	Does the patient need more assistance with transport (either private or public)? (If the patient has difficulties due only to physical problems, e.g bad leg,	tick 'no')		
-				

(To get a total score, add the number of items answered 'no', 'don't know' or 'N/A'

Total score (out of 6)

 $\ \ \, \textbf{If patient scores 0-3, cognitive impairment is indicated. Conduct standard investigations.}$

© University of New South Wales as represented by the Dementia Collaborative Research Centre - Assessment and Better Care, Brodaty et al, *JAGS* 2002; 50:530-534

Mini-COG (Patient)

Quick facts:

- •Time to administer: 2-3 min, no more than 5 min
- Domains tested: visuospatial, executive, memory
- Accessibility: Available in multiple languages
- Not adjusted for a patient's education level

Mini-Cog makeup:

- 1.Recall test of three words
- 2.Clock drawing test

Scoring: There are a total of 5 points. Less than 3 points is abnormal.

0-2 indicates a high likelihood of dementia.

3-5 indicates a low likelihood of dementia.



Mini-Cog¹

Instructions for Administration & Scoring

ID: Date:		

Step 1: Three Word Registration

Look directly at person and say."Please listen carefully. I am going to say three words that I want you to repeat back to me now and try to remember. The words are [select a list of words from the versions below]. Please say them for me now If the person is unable to repeat the words after three attempts. move on to Step 2 (clock drawing).

The following and other word lists have been used in one or more clinical studies." For repeated administrations. use of an alternative word list is recommended.

Version 1 Version 2 Version 3 Version 4 Version 5 Version 6

Banana Leader Village River Captain Daughter Sunrise Season Kitchen Nation Garden Heaven Chair Table Baby Finger Picture Mountain

Step 2: Clock Drawing

Say: 'Next I want you to draw a clock for me. First put in all of the numbers where they go: When that is completed say; 'Now, set the hands to 10 past h."

Use preprinted circle (see next page) for this exercise. Repeat instructions as needed as this is **not** a memory test Move to Step 3 if the clock is not complete within three minutes.

Step 3: Three Word Recall

Ask the person to recall the three words you stated in Step 1. Say: 'What were the three words I asked you to remember? Record the word list version number and the person's answers below.

Word List	Version:	Person'	s A	Answers
-----------	----------	---------	-----	---------

Scoring


Word Recall: _ (0-3 points)	1 point for each word spontaneously recalled without cueing.
Clock Draw: _ (0 or 2 points)	Normal dock - 2 points. A normal dock has all numbers placed in the correct sequence and approximately correct position (e.g. 12. 3. 6 and 9 are in anchor positions) with no missing or duplicate numbers. Hands are pointing to [hell and 2 (RIO). Hand length is not scored. Inability or refusal to draw a clock (abnormal) - 0 points.
Total Score: _ (0-5 points)but	Total score - Word Recall score • Clock Draw score. A cut point of <3 on the Mini-Cog" has been validated for dementia screening, many individuals with clinically meaningful cognitive impairment will score higher. When greater sensitivity is desired. a cut point of <4 is recommended as it may indicate a need for further evaluation of cognitive status.

Mini.Cog" L. S. Borson.All rights reserved. Reprinted with permission of the author solely for clinical and educational purposes. May not be modified or used for commercial. marketing. or research purposes without permission of the author (soobuwedu).

v. 01.1916

Clock Drawing

ID:	Date:
ID	Date.



References

1.8orson S. Scanlan JM. Chen PJ et al. The Mini-Cog as a screen for dementia: Validation in a population-based sample. J Am Geriatr Soc 2003:51:1451:1454.

Borson S. Scanlan JM. Watanabe Jet al. Improving identification of cognitive impairment in primary tare. Int J Geriatr Psychiatry 2006:21: 349-355.

- 3. Lessig M. Scanlan J et al.Time that tells: Critical clock-drawing errors for dementia screening. Int Psychogeriatr. 2008 June: 20(3): 459·470.
- 1. Tsoi K. Chan Jet a I. Cognitive tests to detect dementia: A systematic review and metaanalysis. JAMA Intern Med. 2015: 61-69.
- McCarten J. Anderson Petal. Screening for cognitive impairment in an elderly veteran population: Acceptability and results using different versions of the Mini-Cog. J Am Geriatr Soc 20n:59: 30q-213.
- 3. McCarten J. Anderson Petal. Finding dementia In primary care: The results of a clinical demonstration project. J Am Geriatr Soc 2012:60:210-217.
- 4. Scanlan J & Borson S. The Mini-Cog: Receiver operating characteristics with the expert and naive raters. Int J Geriatr Psychiatry 2005:16:216-222.

Mini.Cog' v S. Borson.All rights reserved. Reprinted with permission of the author solely for clinical and educational purposes. May not be modified or used for commercial. marketing. or research purposes without permission of the author (sools.puwedu).

Table of Commonly Asked ADLsZIADLs

Activities of Daily Living

Activities of Daily Living

Bathing

Dressing

Transferring from bed

to chair

ToiLeting

Grooming

Feeding oneself

Instrumental Activities of Daily Living

Using the telephono

Preparing meals

Managing housenoLd

finances

Taking medications

Doing Laundry

Doing housework

Shopping

Managing

transportation

3. Establish and document a patient's support person and/or a health care agent.

Many people may be involved in a person's care to different degrees and for different purposes. There are three roles to define that are involved in the CHA process:

- an informant- can give you information
- a support person- someone who helps with care
- a health care agent- someone who has legal authority if that person is unable to make their own decisions



"Do you have anyone who is available to speak with me and who knows you well?"

Case Study: Mrs. Pérez and Ana

TAKE A BRIEF HISTORY:

Mrs. Pérez—72 years old—reports progressive difficulty remembering what she has to do in a day.

She forgot to pick up her grandson at school and sometimes gets "lost in time" and forgets what she's doing in the middle of a task.

She brings her daughter Ana to the visit today.

You decide to use the GP-COG for both Mrs. Perez and Ana, since it's web-based and automatically scores.





Screening Test: GPCOG

RESULTS OF THE GPCOG TEST

Testing date: 3 June 2022

Patient name: Mrs. Perez

Date of birth: 1 March 1950



With her challenges in recall skills and the clock, Mrs. Perez score of 4 indicates cognitive impairment

The score for this section is: 4 (+)





More information is required. Proceed with informant interview (/index/informant-interview).

GP-COG Function: Informant Interview with Ana

INFORMANT INTERVIEW

You review Mrs. Perez's

ADLs and IADLs, and sepents she can perform all activities independently.

The score for this section is: 4

ASSESSMENT

No significant cognitive impairment. Further testing not necessary. Follow-up in 12 months.



What is the result of the cognitive and functional tests for Mrs. Pérez?

- 1. Negative for cognitive decline, negative for functional decline
- 2. Negative for cognitive decline, positive for functional decline
- 3. Positive for cognitive decline, negative for functional decline
- 4. Positive for cognitive decline, positive for functional decline



Interpret the results

Mrs. Pérez is POSITIVE on her Cognitive Health Assessment.

- 1. Positive symptoms
- 2. Positive (abnormal) on cognitive screening tool
- 3. Negative (normal) on functional screening tool
- 4. (Documented care partner information)

Disclose this to her.



Mrs. Pérez: Documentation

In your note:

- History: Reported more forgetting to pick up her grandson
- Exam: GP-COG part 1 4/9 (positive) and part 2 4/6 (negative)
 - Informant: daughter, Ana
 - Interpretation: negative screen for dementia at this time
- Support system: Ana last name, daughter, contact information
 - Does not have health care agent designated
- GP-COG result disclosed to patient, plan for annual screening and brain health plan

Disclosure

You inform Mrs. Pérez that her GP-COG was positive for possible decline in her cognitive abilities, but her ability to do her daily activities is not affected. You want to continue to do some evaluation though given that she has some symptoms (her screen is positive).

1. "Mrs. Pérez, what is your understanding of the assessment so far?"

1. "What questions do either of you have so far about the assessment?"



Documentation

What Documentation Is Required for Billing?

Billing can occur for 1494F if the provider documents all of the following:

- The screening tool or tools that were used (at least one cognitive screening tool is required in order to bill this code).
- That the completed assessments were reviewed by the provider.
- The results of the assessment(s).
- The interpretation of the results.
- That the results were discussed with the member, family, or informant, and any appropriate actions were taken.



After a positive cognitive health assessment: Start a brain health plan




Make sure vision and hearing assessments are up to date and, if impairments are present, correct them accordingly.



Review medication for cognitive side effects and reduce as many of these as you can in dose or frequency, and preferably stop them. Some common medications with cognitive side effects are listed below.



Encourage social and physical activity.



Continue to address blood pressure and diabetes management goals.



After a positive cognitive health assessment: Plan for a visit to start the next steps in the workup





Tip: Utilize Your Team

Different members of the team can contribute to the CHA. For example, the

front desk staff might notice that a patient is missing more appointments, or

a medical assistant might alert a provider that a patient has been repeating

themselves or coming to appointments on the wrong day.



Thank You!

For more information please contact:

ucigeriatrics@hs .uci.edu







Questions



How to Claim Continuing Medical Education (CME) Credit?

Step 1. Please complete our evaluation survey using the link provided in the chat and a follow-up email after the webinar. For this activity, we provide CME, CAMFT credits. Please select the correct link based on the credit type you are claiming.

You can also scan a QR code on this slide to claim credit NOW!

Step 2. Upon completing the evaluation survey, please scan a QR code or link to claim credit.

- Use your phone camera to scan a QR code and tap the notification to open the link associated with the CME portal.
- Enter you first name, last name, profession, and claim 1 CE
 credit for the webinar.





UC Irvine ECHO Series

Age Friendly Geriatrics Education

- Every other Wednesday and every Friday at the lunch hour (12-1PM PST)
- Multitude of topics in relation to Geriatrics, with a new topic every other Wednesday
- Open to all specialties, Nurse
 Practitioners, Physician Assistants, etc.
- Free CME/CE/CEU's per session!!!

Scan the QR code on flyer to register for our series



Age-Friendly

Geriatrics Education

Learn how to provide better care for older adults by applying the 4Ms of geriatric care: Mind, Medication, Mobility, & What Matters Most. Build geriatric competencies for better patient outcomes. Improve dementia care and cognitive health screening.



Why Attend?

- Interactive Real-Time Consultative Discussion with Board Certified Geriatrics Experts Across the UC System
- · Develop a Specialty in Geriatrics
- Receive FREE AMA PRA Category 1 Credit™ per session

Target Audience



All Health Care Professionals Seeking to Improve the Care of Older Adults





JOIN US ON ZOOM

Every Other Wednesday and Every Friday 12:00-1:00PM PST



Scan QR Code to Register



