

| HEDIS Trends in D-SNP (HEDIS MY 2021-MY 2023) | | | | | | | |
|--|--|-------------------------|-------------------------|-------------------------|--|--|--|
| Acronyms | Measures | HEDIS 2022 (MY 2021) | HEDIS 2023 (MY 2022) | HEDIS 2024 (MY 2023) | | | |
| AAB | Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis - Total | 46.60% | 38.93% | 31.30% | | | |
| AAP | Adults Access to Preventive/Ambulatory Health Services (Ages 65 and older) | 87.85% | 88.38% | 88.57% | | | |
| AMM | Antidepressant Medication Management – Acute Phase | 75.65% | 77.66% | 77.15% | | | |
| AMM | Antidepressant Medication Management – Continuous Phase | 57.86% | 54.71% | 57.99% | | | |
| BCS-E | Breast Cancer Screening | 62.84% | 68.15% | 66.89% | | | |
| BPD | Blood Pressure Control (<140/90) for Patients With Diabetes | 64.09% | 73.96% | 76.01% | | | |
| СВР | Controlling High Blood Pressure | 61.81% | 70.92% | 74.71% | | | |
| COL-E | Colorectal Cancer Screening | N/A | 60.12% | 62.78% | | | |
| DDE | Potentially Harmful Drug-Disease Interactions in the Elderly – Total (lower rate is better) | 34.68% | 34.13% | 36.12% | | | |
| EED | Eye Exam for Patients With Diabetes | 68.83% | 78.91% | 74.12% | | | |
| FUA | Follow-up After Emergency Department Visit for Alcohol and other Drug Abuse or Dependence – 7-day rate | 9.38% | 26.76% | 23.53% | | | |
| FUH | Follow-Up after Hospitalization for Mental Illness –within 30-days | 63.43% | 58.33% | 61.11% | | | |
| FUH | Follow-Up after Hospitalization for Mental Illness -within 7-days | 36.57% | 34.62% | 46.03% | | | |
| FUM | Follow-up After Emergency Department Visit for Mental Illness – 30-day rate | 52.00% | 43.75% | 48.84% | | | |
| FUM | Follow-up After Emergency Department Visit for Mental Illness - 7-day rate | 41.33% | 31.25% | 38.37% | | | |



| HEDIS Trends in D-SNP (HEDIS MY 2021-MY 2023) | | | | | | | |
|--|---|-------------------------|-------------------------|-------------------------|--|--|--|
| Acronyms | Measures | HEDIS 2022 (MY 2021) | HEDIS 2023 (MY 2022) | HEDIS 2024 (MY 2023) | | | |
| HBD | Hemoglobin A1c Control for Patients With Diabetes — HbA1c Control (<8%) | 63.84% | 66.15% | 62.80% | | | |
| HBD | Comprehensive Diabetes Care - Poor HbA1c Control >9% (lower rates is better) | 24.94% | 23.44% | 26.68% | | | |
| HDO | Use of Opioids at High Dosage (lower rates is better) | 3.14% | 3.24% | 1.84% | | | |
| IET | Initiation and Engagement of Alcohol and Other Drug Dependence Treatment - Engagement | 6.68% | 8.49% | 7.57% | | | |
| IET | Initiation and Engagement of Alcohol and Other Drug Dependence Treatment – Initiation | 55.26% | 50.84% | 50.91% | | | |
| KED | Kidney Health Evaluation for Patients With Diabetes | N/A | 64.26% | 65.36% | | | |
| OMW | Osteoporosis Management in Women Who Had a Fracture | 39.34% | 56.25% | 44.64% | | | |
| OSW | Osteoporosis Screening in Older Women | 37.42% | 44.41% | 48.39% | | | |
| PCE | Pharmacotherapy Management of COPD Exacerbation- Bronchodilators | 92.14% | 89.60% | 83.80% | | | |
| PCE | Pharmacotherapy Management of COPD Exacerbation -Systemic Corticosteroid | 71.43% | 79.20% | 68.31% | | | |
| PSA | Non-Recommended PSA-Based Screening in Older Men (lower rates is better) | 34.33% | 36.40% | 33.09% | | | |
| SMD | Diabetes Monitoring for People With Diabetes and Schizophrenia | 78.10% | 71.88% | 81.44% | | | |
| SPC | Statin Therapy for Patients with Cardiovascular Disease - Therapy | 80.28% | 82.55% | 85.76% | | | |
| SPC | Statin Therapy for Patients with Cardiovascular Disease – Adherence | 76.29% | 75.43% | 84.84% | | | |
| SPD | Statin Therapy for Patients with Diabetes - Therapy | 82.10% | 82.14% | 82.51% | | | |



| HEDIS Trends in D-SNP (HEDIS MY 2021-MY 2023) | | | | | | |
|--|---|-------------------------|-------------------------|-------------------------|--|--|
| Acronyms | Measures | HEDIS 2022 (MY 2021) | HEDIS 2023 (MY 2022) | HEDIS 2024 (MY 2023) | | |
| SPD | Statin Therapy for Patients with Diabetes – Adherence | 74.25% | 74.75% | 82.35% | | |
| SSD | Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications | 74.95% | 78.23% | 85.86% | | |
| TRC | Transitions of Care - Receipt of Discharge information: | 3.89% | 6.81% | 3.16% | | |
| TRC | Transitions of Care - Patient engagement after inpatient discharge | 81.27% | 69.34% | 75.67% | | |
| TRC | Transitions of Care - Notification of Inpatient Admission | 5.11% | 10.71% | 3.89% | | |
| TRC | Transitions of Care - Medication reconciliation post discharge | 39.42% | 42.34% | 32.12% | | |
| UOP | Use of Opioids from Multiple Providers - Multiple Prescribers and Multiple Pharmacies Rates (lower rates is better) | 0.88% | 0.65% | 1.04% | | |